Journal of Applied Biology & Biotechnology Vol. 13(1), pp. 75-82, Jan-Feb, 2025

Available online at http://www.jabonline.in

DOI: 10.7324/JABB.2024.159375



## Gumghatti-based composite coating improves postharvest quality and nutritional value of black plums

T. V. Ramana Rao<sup>1</sup>\*, Pinal K. Dave<sup>2</sup>, Ananya Payal<sup>1</sup>, Jay B. Pandya<sup>2</sup>, Prakash R. Patel<sup>2</sup>, Neha Pareek<sup>2</sup>

#### ARTICLE INFO

Article history:

Received on: May 30, 2024 Accepted on: September 25, 2024 Available online: November 15, 2024

## Key words:

Black plum, Composite coating, Nutritional quality, Chamomile oil, Jojoba oil.

#### **ABSTRACT**

Functional materials, along with plasticizers, surfactants, and other additives, dissolved in the right proportions in : various solvents form edible coatings play a significant role in the extension of fresh produce's shelf-life. The aim of the study was to improve the quality of black plum during storage ( $Prunus\ salicina\ Lindl.$ ) using gumghatti and lipid-based composite edible coatings. Different concentrations of gumghatti in combination with oils (coconut oil, jojoba oil, and chamomile oil) were coated onto plum fruits and stored at a temperature of  $24\pm5^{\circ}C$  for 16 days. The physicochemical and biochemical changes occurring during the ripening of treated and untreated black plums were analyzed. Untreated plum fruits exhibited higher activity of cell wall softening enzymes compared with those treated with gumghatti alone and other formulations. Fruits treated with gumghatti and chamomile oil retained firmness better (6 lb) throughout the storage period. Treatment of plums with gumghatti in combination with chamomile oil and jojoba oil showed a significant impact on the retention of bioactive compounds like antioxidants, ascorbic acid, phenols, and anthocyanins. The findings offer suggestions for an economical and convenient alternative to maintain the keeping quality of black plum fruits stored at room temperature.

## 1. INTRODUCTION

Plums (*Prunus salicina* Lindl.), popularly known as stone fruits, belong to the Rosaceae family, which are nutritious in terms of minerals, vitamins, phytochemicals (i.e., phenolic acids, anthocyanins, and carotenoids), and also low in calories [1]. However, plums have a very low shelf life of 3–4 days, further resulting in the postharvest losses due to fruit softening, chilling injuries, and browning issues [2]. To increase their shelf life after harvest and maintain their quality, it is recommended to store plums at 0°C [3].

It is essential to be cautious with cold storage as it can lead to softening and chilling injuries in most tested plum cultivars, as observed by Vangdal *et al.* [4]. To address this, researchers have explored various postharvest technologies like the application of methyl cyclopropane

\*Corresponding Author:

T. V. Ramana Rao, Department of Food Technology, VIGNAN's Foundation for Science, Technology & Research, (Deemed to be University), Guntur, Andhra Pradesh, India

E-mail: drtvrr ft@vignan.ac.in

(1-MCP) [5], modified atmosphere packaging [6], and edible coatings [7,8] to preserve the quality and extend the storage life of plums.

Application of the edible coatings has become a promising approach to enhance the shelf life of various fresh produces. This can be achieved by reducing food-borne pathogens, maintaining firmness, and preventing weight loss during storage for up to a month [9]. Edible coatings can be made from proteins, polysaccharides, lipids, resins, or combinations of these materials. In addition to these, functional ingredients like essential oils, antioxidants, and flavors can also be incorporated aiming toward the better quality, stability, safety, and functionality of food during processing, handling, and storage [10]. Selection of coating material depends on factors such as water solubility, hydrophilicity, hydrophobicity, ease of formation, and sensory properties.

One such coating material is gumghatti, also known as Indian gum, which is a complex nonstarch polysaccharide widely utilized in the food and pharmaceutical industries due to its excellent emulsification and thickening properties [11]. Combining gumghatti with various edible/essential oils having antibacterial and antifungal properties has shown promising results. For example, an edible coating formulated using gumghatti and clove oil emulsion enhanced the shelf life of papaya, along with preserving its quality throughout the storage

<sup>&</sup>lt;sup>1</sup>Department of Food Technology, Vignan's Foundation for Science, Technology and Research, Guntur, Andhra Pradesh, India.

<sup>&</sup>lt;sup>2</sup>Department of Biosciences, Sardar Patel University, Vallabh Vidyanagar, Gujarat, India.

period. This approach not only conserves energy but also minimizes the risk of cold storage disorders during fruit preservation [11]. Therefore, the objective of the present research is to investigate the shelf life enhancement of plums during the postharvest storage by retaining their physicochemical characteristics using an affordable and eco-friendly method of applying edible coatings (i.e., gumghatti combination with coconut oil, jojoba oil, and chamomile oil) at room temperature.

### 2. MATERIALS AND METHODS

### 2.1. Sample Selection and Postharvest Treatment

The initial step involved in the study was the selection of black plum fruits that were of uniform maturity, shape, and size, with no physical damage. Prior to the coating process, fruits were washed carefully with water, followed by disinfection using a 2% sodium hypochlorite solution. Finally, the fruits were kept to air-dry at ambient temperature.

The plum fruits were graded into five sets, each set consisting of 15 fruits. The sets T1, T2, T3, and T4 were subjected to distinct formulations of edible coating emulsions, while the fifth set served as the control (C). After the application of coatings on the fruits, they were kept for surface drying at room temperature. Subsequently, the treated and control fruits were stored at a temperature of  $24 \pm 5^{\circ}\text{C}$  and relative humidity ranging from 44% to 62%. Throughout the 16-day storage period, the stored fruits were assessed for their physicochemical, biochemical, and microbial properties at intervals of 0, 4, 8, 12, and 16 days.

## 2.2. Viscosity Measurement of the Coatings

The rheological analysis aimed to investigate the viscosity of various concentrations of gumghatti (0.3%, 0.6%, and 0.9%) both individually and in conjunction with coconut oil, jojoba oil, and chamomile oil (0.2 and 0.3%). After analyzing the initial findings [Table 1], the study focused on intermediate viscous concentrations of the different coatings. Optimized concentrations of gumghatti were employed alongside three oils: coconut oil, jojoba oil, and chamomile oil. According to rheological analysis, gumghatti alone (0.6%) or in conjunction with oils, like gumghatti (0.6%) + coconut oil (0.3%)

**Table 1:** Viscosities of different formulations of gumghatti and combination of oils like coconut oil, jojoba oil, and chamomile oil.

or ons inte coe	onat on, jojooa on, and enamonnie on.	
Treatments	Sample	Viscosity (mPa)
TA1	0.3% gum-ghatti	11.782
TA2	0.6% gum-ghatti	6.461
TA3	0.9% gum-ghatti	5.672
TA4	0.6% gum-ghatti + $0.2%$ coconut oil	2.707
TA5	0.6% gum-ghatti + $0.3%$ coconut oil	2.930
TA6	0.9% gum-ghatti + 0.3 % coconut oil	3.744
TA7	0.6% gum-ghatti + $0.2%$ jojoba oil	3.667
TA8	0.6% gum-ghatti $+$ $0.3%$ jojoba oil	25.06
TA9	0.9% gum-ghatti + $0.3%$ jojoba oil	16.24
TA10	0.6% gum-ghatti + 0.2% chamomile oil	2.679
TA11	0.6% gum-ghatti + 0.3% chamomile oil	3.14
TA12	0.9% gum-ghatti + 0.3% chamomile oil	6.1543

and gumghatti (0.6%) + chamomile oil (0.3%), exhibited moderate viscosity, making them the chosen applicants for the study. Table 2 presents the concentrations and viscosities of the coatings.

## 2.3. Preparation of Coating Formulations

T1: 0.6% gumghatti

T2: 0.6% gumghatti + 0.3% coconut

T3: 0.9% gumghatti + 0.3% jojoba oil

T4: 0.6% gumghatti + 0.3% chamomile oil

The abovementioned mixture was filtered to get better purity, and distilled water was added to obtain a final volume of 100 ml. To reduce surface tension, Tween-80 was incorporated into all the coatings. A magnetic stirrer was used to make a uniform solution. After the addition of 0.75% glycerol, the plasticizer component solution was again stirred for 10 min. The resulting formulations were then applied to four sets of fruits, with the fifth set serving as a control to compare the ripening process between coated and uncoated fruits.

## 2.4. Physicochemical and Biochemical Analyses

The efficacy of the edible coatings was assessed by evaluating various physicochemical parameters:

The percentage of decay was calculated using the formula

Decay % = ((Initial no. of fruits - Final no. of fruits) / Final no. of fruits) x 100

The percentage of weight reduction was assessed using the formula

Weight loss % = ((Initial weight of fruits - Final weight of fruits) / Initial weight of fruits) x 100

The shelf life of the coated fruits was evaluated following the below methods described by Mondal [12]. The pH and total soluble solids (%) (TSS) were measured using the AOAC method [13]. The total sugar content was analyzed using Sadasivam and Manickam's method [14]. The browning index was calculated as described by Eissa [15]. Anthocyanins and flavanols were measured using Lees and Francis's method [16] to assess the presence of antioxidants. The total phenolic content and total antioxidant activity of the samples were evaluated by the methods described by Larrauri *et al.* [17] and McDonald *et al.* [18], respectively. Ascorbic acid, carotene, and lycopene content were analyzed by the methods described by Roe and Keather [19] and Wang *et al.* [20].

**Table 2:** Treatments considered for the coating application on black plum fruits.

Treatment	Viscosity (mPa)
T1	6.461 (TA2)
T2	2.930 (TA5)
T3	16.24 (TA5)
T4	3.14 (TA11)

T1: Gum-ghatti 0.6%; T2: Gum-ghatti (0.6%) + coconut oil (0.3%); T3: Gum-ghatti (0.9%)+ jojoba oil (0.3%) and T4: Gum-ghatti (0.6%) + chamomile oil (0.3%).

To observe the efficacy of the edible coatings, the degrading activity of cell wall enzymes, like pectate lyase (PL) and polygalacturonase (PG), was determined using the methods of Ouattara *et al.* [21], Pathak and Sanwal [22], and Eissa [15], respectively. The activity of polyphenol oxidase enzyme responsible for browning was also evaluated using Xing *et al.* [23] method.

## 2.5. Statistical Analysis

All the analyses were performed in triplicates (n=3) and represented as mean  $\pm$  SEM. Following the methodology described by Bliss [24], the data from the present study were put through Duncan's multiple range test to assess the significant difference between the obtained parameters (p < 0.05). To detect the significant differences among all the treatments and control set, the overall least significant difference ( $p \le 0.05$ ) was also calculated.

### 3. RESULTS AND DISCUSSION

## 3.1. Weight Loss and Decay

A significant ( $p \le 0.05$ ) increase in weight loss percentage was observed, which can be attributed to the increased respiration activity following harvest. However, the treated fruits exhibited lower weight loss compared with the control group. Among the treated fruits, those treated with jojoba oil showed significant changes in weight loss percentage ( $p \le 0.05$ ). For instance, after 8 days of storage, the weight loss in gumghatti + jojoba oil (T3) was 9.30%, while the control group recorded a weight loss of 24.41%. Additionally, the decay percentage was found to be lower in the treated fruits, with only 20% decay observed in the T4 treatment, as opposed to the 40% decay in the control group after 8 days of storage. The reduction in weight loss and decay has been reported earlier in papaya fruit with gumghatti 3% and clove oil 0.1% [11].

Moreover, Rojas-Argudo elucidated that the incorporation of lipids, along with the polysaccharide coatings, can enhance the water barrier nature of the coatings [25]. In line with the above study, edible coatings in the current research also reported better effectiveness due to the addition of a lipid component, chamomile oil with gumghatti, indicating the maintenance of the hydrophilic–hydrophobic balance, which results in the reduction in water loss from the fruit. The above findings were also in accordance with Kittur *et al.*, who reported the reduction in weight loss when polysaccharide-based composite

coatings were applied on banana fruits compared with uncoated ones [26]. In another study, it was also noticed that the coatings incorporated with composite oil were able to retard the ethylene emission in pineapple and reduce the weight loss during storage [27].

#### 3.2. Firmness

An increase in the storage duration lead to a decrease in the firmness of the fruits significantly ( $p \le 0.05$ ). Nevertheless, fruits treated with a combination of gumghatti and chamomile oil (T4) maintained their firmness during the entire storage period compared with the control and other formulations [Table 3]. Retention of firmness in the coated fruits could be attributed to the reduction in PG and pectinesterase activities that are responsible for breaking of bonds in the pectin present in the cell wall, thus leading to the disintegration of insoluble protopectins to more soluble pectic acid and pectins [28]. Higher percentage of carbon dioxide and lower concentrations of oxygen also reduce the activities of these enzymes, allowing the coated fruits to retain the firmness better [29].

Therefore, the results of this study show a lot of resemblance to the findings by Mahfoudhi and Hamdi [30], who noticed that the maximum firmness in cherry fruit was maintained by coatings with gum arabic and almond gum and also with the reports of Pandey *et al.*, confirming the role of composite edible coatings in retarding ethylene emission and enhancing texture, thereby preserving the quality of fruits [31].

## 3.3. pH and TSS

During storage, initially pH was found to be  $4.043 \pm 0.015$  and a slight decrease was observed in all fruits during the storage. T3 (3.306  $\pm$  0.005-16th day) and T4 (3.273  $\pm$  0.005-16th day) exhibited a slower decrease in pH compared with other groups (T1-3.223  $\pm$  0.005-12th day, T2-3.910  $\pm$  0.020-16th day, C-3.456  $\pm$  0.005-8th day); however, there were no notable variations in pH between the treated and control fruits. On the contrary, both control and treated fruits exhibited an increase in TSS until the eighth day, after which it gradually declined in coated fruits [Table 3]. A large amount of polysaccharides (e.g., pectin and cellulose) present in the cell wall will be digested by the degrading enzymes, leading to an increase in the soluble solid content during the process of ripening; however, they are utilized for the respiration process in the later stages [32]. The changes in TSS were significantly slower in coated fruits than in control ( $p \le 0.05$ ), proving the efficacy of edible coatings as selective barriers to CO<sub>2</sub> and O<sub>2</sub>, thus modifying

	Table 3: Change	s in firmness and total	l soluble solids of black	plum fruits during s	torage at $25 \pm 1$ °C.
--	-----------------	-------------------------	---------------------------	----------------------	--------------------------

	Day 0	Day 4	Day 8	Day 12	Day 16
Treatments			Firmness (lb)		
T1	16.0	7.0	6.0	6.0	-
T2	16.0	10.0	9.0	5.0	-
T3	16.0	8.5	6.0	4.0	-
T4	16.0	14.0	10.0	5.0	6.0
C	16.0	8.0	3.0	-	-
Treatments			Total soluble solids (%)		
T1	$0.5\pm0^{\mathrm{a}}$	$1.43 \pm 0.06^{\circ}$	$1.37\pm0.12^{\rm d}$	$1.20\pm0.00^{\text{b}}$	-
T2	$0.5\pm0^{\mathrm{a}}$	$1.03\pm0.06^a$	$1.13 \pm 0.06^{b}$	$1.20 \pm 0.00^{\rm b}$	$1.20\pm0.00^{\rm b}$
T3	$0.5\pm0^{\mathrm{a}}$	$1.47\pm0.06^{\rm d}$	$1.10\pm0.00^{\rm a}$	$1.10\pm0.00^{\rm a}$	$0.83\pm0.06^{\rm a}$
T4	$0.5\pm0^{\mathrm{a}}$	$1.37\pm0.06^{\rm b}$	$1.30 \pm 0.00^{\circ}$	$1.17\pm0.06^{\rm c}$	$1.33 \pm 0.06^{\circ}$
C	$0.5\pm0^{\mathrm{a}}$	$1.50\pm0.00^{\rm e}$	$1.37\pm0.06^{\text{e}}$	-	-

Values are mean  $\pm$  standard deviation, n = 3. Values within treatments with different letters (a–d) in a column differ significantly ( $p \le 0.05$ ) with values from higher to lower. T1: Gumghatti 0.6%; T2: Gum-ghatti (0.6%) coconut oil (0.3%); T3: Gum-ghatti (0.9%) + jojoba oil (0.3%) and T4: Gum-ghatti (0.6%) + chamomile oil (0.3%) and C: control.

Table 4: Changes in anthocyanins, flavanols, and lycopene content of black plum fruits during storage at 25 ± 1°C.

	Day 0	Day 4	Day 8	Day 12	<b>Day 16</b>
Treatments			Anthocynins (μg g <sup>-1</sup> FW)		
T1	$1.711 \pm 0.079^{a}$	$2.810 \pm 0.003^{\rm d}$	$2.799 \pm 0.016^{c}$	$3.679 \pm 0.039^{\rm d}$	-
T2	$1.711 \pm 0.079^{a}$	$2.551 \pm 0.013^{b}$	$2.665 \pm 0.008^{\rm b}$	$2.939 \pm 0.024^{c}$	$1.218 \pm 0.063^{c}$
T3	$1.711 \pm 0.079^{a}$	$2.745 \pm 0.026^{c}$	$1.363 \pm 0.009^{\rm a}$	$1.499 \pm 0.005^{\rm a}$	$1.198 \pm 0.023^{\rm b}$
T4	$1.711 \pm 0.079^{a}$	$2.515 \pm 0.015^a$	$3.236 \pm 0.003^{\rm d}$	$2.847 \pm 0.000^{\rm b}$	$1.118 \pm 0.005^{\rm a}$
C	$1.711 \pm 0.079^{a}$	$3.295 \pm 0.008^{e}$	$3.662 \pm 0.006^{e}$	-	-
Treatments			Flavanols (µg g <sup>-1</sup> FW)		
T1	$3.089 \pm 0.019^{\rm e}$	$2.578 \pm 0.012^{\rm d}$	$3.250 \pm 0.016^{b}$	$6.185 \pm 0.035^{\rm d}$	-
T2	$3.089 \pm 0.019^{e}$	$2.065 \pm 0.013^a$	$4.202 \pm 0.015^{\rm c}$	$5.181 \pm 0.016^{c}$	$2.134\pm0.031^{\text{c}}$
T3	$3.089 \pm 0.019^{\rm e}$	$2.099 \pm 0.018^{b}$	$1.686 \pm 0.003^{\rm a}$	$0.996 \pm 0.003^a$	$2.122 \pm 0.077^{b}$
T4	$3.089 \pm 0.019^{e}$	$2.211 \pm 0.009^{c}$	$4.386 \pm 0.005^{\rm d}$	$3.315 \pm 0.003^{\rm b}$	$1.372 \pm 0.003^{\rm a}$
C	$3.089 \pm 0.019^{\rm e}$	$3.709 \pm 0.005^{\rm e}$	$5.155 \pm 0.003^{\rm e}$	-	-
Treatments			Lycopene (μg g <sup>-1</sup> FW)		
T1	$0.289 \pm 0.001^a$	$0.526 \pm 0.001^{e}$	$0.846 \pm 0.001^{\rm b}$	$1.407 \pm 0.010^{\rm d}$	-
T2	$0.289 \pm 0.001^{\rm a}$	$0.520 \pm 0.002^{\rm d}$	$2.484 \pm 0.056^{\rm e}$	$0.834 \pm 0.013^{\rm a}$	$0.814 \pm 0.011^{\rm b}$
T3	$0.289 \pm 0.001^{\rm a}$	$0.464 \pm 0.001^{\circ}$	$0.859 \pm 0.064^{c}$	$0.879 \pm 0.019^{c}$	$0.404 \pm 0.007^a$
T4	$0.289 \pm 0.001^{\rm a}$	$0.308 \pm 0.001^a$	$0.836\pm0.030^a$	$0.862 \pm 0.023^{\rm b}$	$0.940\pm0.007^{\text{c}}$
C	$0.289 \pm 0.001^{\rm a}$	$0.313 \pm 0.001^{b}$	$0.934 \pm 0.020^{\rm d}$	-	-

Values are mean  $\pm$  standard deviation, n = 3. Values within treatments with different letters (a–d) in a column differ significantly (p  $\leq$  0.05) with values from higher to lower. T1: Gunghatti 0.6%; T2: Gunghatti (0.6%)+ coconut oil (0.3%); T3: Gunghatti (0.9%)+ jojoba oil (0.3%) and T4: Gunghatti (0.6%)+ chamomile oil (0.3%) and C: control.

Table 5: Changes in total phenols, ascorbic acid, and total antioxidant activity of black plum fruits during storage at 25 ± 1°C.

	Day 0	Day 4	Day 8	Day 12	Day 16
Treatments		ŗ	Total phenols (mg g <sup>-1</sup> FW	)	
T1	$2.162 \pm 0.007^{e}$	$2.238 \pm 0.005^{\rm d}$	$1.286 \pm 0.009^{\rm e}$	$0.832 \pm 0.004^{\rm d}$	-
T2	$2.162 \pm 0.007^{\rm e}$	$2.529 \pm 0.006^{e}$	$0.465 \pm 0.002^a$	$0.952 \pm 0.004^{b}$	$0.558 \pm 0.002^{\rm a}$
T3	$2.162 \pm 0.007^{\rm e}$	$1.329 \pm 0.002^{b}$	$0.516 \pm 0.008^{b}$	$0.251 \pm 0.001^a$	$0.805 \pm 0.001^{\rm b}$
T4	$2.162 \pm 0.007^{e}$	$2.214 \pm 0.005^{c}$	$0.736 \pm 0.013^{\rm d}$	$0.939 \pm 0.004^c$	$0.952 \pm 0.004^{c}$
C	$2.162 \pm 0.007^{\rm e}$	$1.270 \pm 0.013^{\rm a}$	$0.595 \pm 0.002^{c}$	-	-
Treatments		As	corbic acid (mg 100 g <sup>-1</sup> F	W)	
T1	$37.50 \pm 0.00^{\rm e}$	$30.42 \pm 0.01^{e}$	$17.71 \pm 0.01^{a}$	$51.88 \pm 0.00^{d}$	-
T2	$37.50 \pm 0.00^{\rm e}$	$15.42 \pm 0.00^{a}$	$50.21 \pm 0.03^{e}$	$32.50 \pm 0.01^{b}$	$23.75 \pm 0.01^{\circ}$
T3	$37.50 \pm 0.00^{e}$	$18.75\pm0.00^{\rm c}$	$22.29 \pm 0.01^{\rm b}$	$13.75\pm0.01^a$	$18.54\pm0.01^{\mathrm{a}}$
T4	$37.50 \pm 0.00^{e}$	$16.67 \pm 0.00^{\rm b}$	$35.00\pm0.01^{\rm d}$	$43.13 \pm 0.02^{c}$	$20.21 \pm 0.01^{\rm b}$
C	$37.50 \pm 0.00^{\rm e}$	$27.29\pm0.00^{\rm d}$	$28.96\pm0.00^{\rm c}$	-	-
Treatments		Tota	l antioxidant activity (%/	FW)	
T1	$94.554 \pm 0.000^{\rm a}$	$93.812 \pm 0.000^{b}$	$90.890 \pm 0.001^{e}$	$94.015 \pm 0.002^{\circ}$	-
T2	$94.554 \pm 0.000^{a}$	$94.389 \pm 0.001^{\circ}$	$93.240 \pm 0.001^{b}$	$93.629 \pm 0.005^{b}$	$87.387 \pm 0.017^{a}$
T3	$94.554 \pm 0.000^{\mathrm{a}}$	$94.554 \pm 0.001^{\circ}$	$95.890 \pm 0.001^{c}$	$73.488 \pm 0.014^{\rm a}$	$90.927 \pm 0.006^{b}$
T4	$94.554 \pm 0.000^{\mathrm{a}}$	$95.132 \pm 0.001^{\rm d}$	$99.947 \pm 0.001^{\rm d}$	$94.723 \pm 0.001^{\circ}$	$93.308 \pm 0.001^{c}$
C	$94.554 \pm 0.000^{\rm a}$	$80.000 \pm 0.000^{\rm a}$	$73.230 \pm 0.001^{\rm a}$	-	-

Values are mean  $\pm$ standard deviation, n = 3. Values within treatments with different letters (a–e) in a column differ significantly ( $P \le 0.05$ ) with values from higher to lower. T1: Gumghatti 0.6%; T2: Gum-ghatti (0.6% coconut oil (0.3%); T3: Gum-ghatti (0.9%) + jojoba oil (0.3%) and T4: Gum-ghatti (0.6%) + chamomile oil (0.3%) and C: control.

the internal atmosphere of the fruit followed by slowing down the rate of respiration [33].

## 3.4. Anthocyanin and Flavanols

Throughout the study period, the anthocyanin content of control fruit exhibited a gradual increase. However, in the case of treated black plum fruits, the increase in anthocyanin content was slower and sustained as the ripening was delayed compared with the control group [Table 4] that led to the greater anthocyanin content in treated fruits by the end of storage. These findings align similarly with the results obtained from investigation on cherries coated with olive leaves'

extract enriched with combinations of alginate and chitosan were carried out by Zam [34]. Moreover, in black plum fruit, the flavanols initially increased, followed by a declining trend, corroborating the findings from previous research on black plums [35]. Therefore, the stability of anthocyanins during storage duration acting as a scavenger-preventing oxidation of fruit enhances its shelf life.

### 3.5. Lycopene

Fruits coated with (T4) 0.6% gumghatti + 0.3% chamomile oil exhibited less loss of lycopene throughout the storage period ( $p \le 0.05$ ). In comparison, all the treated fruits exhibited lower lycopene

**Table 6:** Changes in polygalactouranase, pectate lyase, and amylase activity of black plum fruits during storage at  $25 \pm 1^{\circ}$ C.

	Day 0	Day 4	Day 8	Day 12	Day 16	
Treatments	Polygalacturonase (U/min/mg protein)					
T1	$0.0217 \pm 0.0033^{\mathrm{a}}$	$0.0330 \pm 0.0040^{\rm b}$	$0.0408 \pm 0.0010^{c}$	$0.0263 \pm 0.0034^{\rm b}$	-	
T2	$0.0217 \pm 0.0033^{\rm a}$	$0.0280 \pm 0.0017^{\rm a}$	$0.0402 \pm 0.0047^{\rm e}$	$0.0632 \pm 0.0157^{\rm d}$	$0.0254 \pm 0.0016^{b}$	
T3	$0.0217 \pm 0.0033^{\rm a}$	$0.0384 \pm 0.0007^{\rm e}$	$0.0566 \pm 0.0044^{b}$	$0.0190 \pm 0.0077^{\rm a}$	$0.0158 \pm 0.0016^{a}$	
T4	$0.0217 \pm 0.0033^{\rm a}$	$0.0340 \pm 0.0012^{\circ}$	$0.0465 \pm 0.0037^{\rm d}$	$0.0563 \pm 0.0102^{\circ}$	$0.0257 \pm 0.0007^{\text{c}}$	
C	$0.0217 \pm 0.0033^{\mathrm{a}}$	$0.0440 \pm 0.0024^{\rm d}$	$0.0504 \pm 0.0004^{a}$	-	-	
Treatments	Pectate lyase (U/min/mg protein)					
T1	$0.0210 \pm 0.0038^{a}$	$0.0277 \pm 0.0003^{\rm b}$	$0.0299 \pm 0.0009^{\rm b}$	$0.0398 \pm 0.0030^{b}$	-	
T2	$0.0210 \pm 0.0038^{\rm a}$	$0.0291 \pm 0.0004^{c}$	$0.0316 \pm 0.0018^{\rm d}$	$0.0423 \pm 0.0022^{c}$	$0.0492 \pm 0.0008^{c}$	
T3	$0.0210 \pm 0.0038^a$	$0.0259 \pm 0.0013^{a}$	$0.0327 \pm 0.0021^{e}$	$0.0391 \pm 0.0004^{\rm a}$	$0.0438 \pm 0.0026^{b}$	
T4	$0.0210 \pm 0.0038^a$	$0.0306 \pm 0.0013^{\rm d}$	$0.0275 \pm 0.0020^{\rm a}$	$0.0425 \pm 0.0021^{\rm d}$	$0.0424 \pm 0.0007^a$	
C	$0.0210 \pm 0.0038^{\mathrm{a}}$	$0.0313 \pm 0.0016^{\rm e}$	$0.0300 \pm 0.0007^{c}$	-	-	
Treatments	Amylase (U/min/mg protein)					
T1	$86.39 \pm 0.319^{a}$	$24.11 \pm 0.18^{\circ}$	$38.23 \pm 0.034^{j}$	$45.05 \pm 0.04^{l}$	-	
T2	$86.39 \pm 0.319^{\mathrm{a}}$	$51.44\pm0.27^{\mathrm{m}}$	$29.13 \pm 0.07^{d}$	$33.77\pm0.05^{\rm g}$	$30.64 \pm 0.02^{e}$	
T3	$86.39 \pm 0.319^{\mathrm{a}}$	$81.43 \pm 0.65^{p}$	$31.78\pm0.04^{\rm f}$	$23.13 \pm 0.03^{b}$	$21.38\pm0.01^{\mathrm{a}}$	
T4	$86.39 \pm 0.319^{\rm a}$	$70.92\pm0.55^{\rm o}$	$34.38\pm0.02^{\rm h}$	$36.64\pm0.08^{\mathrm{i}}$	$40.72 \pm 0.08^k$	
C	$86.39 \pm 0.319^{\rm a}$	$68.24 \pm 0.07^{\rm n}$	$34.67 \pm 0.01^{\rm h}$	-	-	

Values are mean  $\pm$ standard deviation, n = 3. Values within treatments with different letters (a–o) in a column differ significantly ( $P \le 0.05$ ) with values from higher to lower. T1: Gumghatti 0.6%; T2: Gum-ghatti (0.6% coconut oil (0.3%); T3: Gum-ghatti (0.9%)  $\pm$  jojoba oil (0.3%) and T4: Gum-ghatti (0.6%)  $\pm$  chamomile oil (0.3%) and C: control.

Table 7: Changes in browning index and polyphenol oxidase activity of black plum fruits during storage at 25 ± 1°C.

	Day 0	Day 4	Day 8	Day 12	<b>Day 16</b>		
Treatments			Browning index				
T1	$0.134 \pm 0.001^{a}$	$0.212 \pm 0.003^{\rm d}$	$0.224 \pm 0.002^{\rm d}$	$0.280 \pm 0.002^{\rm d}$	-		
T2	$0.134 \pm 0.001^{a}$	$0.165 \pm 0.002^{e}$	$0.251 \pm 0.002^{e}$	$0.150 \pm 0.001^{\circ}$	$0.149 \pm 0.001^{\rm b}$		
T3	$0.134 \pm 0.001^{a}$	$0.221 \pm 0.005^{a}$	$0.127 \pm 0.000^{\rm b}$	$0.096 \pm 0.001^{a}$	$0.145 \pm 0.002^{a}$		
T4	$0.134 \pm 0.001^{a}$	$0.162 \pm 0.006^{b}$	$0.183 \pm 0.003^{\circ}$	$0.124 \pm 0.003^{\rm b}$	$0.162 \pm 0.001^{c}$		
C	$0.134 \pm 0.001^{\rm a}$	$0.145 \pm 0.001^{c}$	$0.217 \pm 0.001^{\rm a}$	-	-		
Treatments		Polyphenol oxidase (U min <sup>-1</sup> mg protein <sup>-1</sup> )					
T1	$0.0217 \pm 0.0033^{\rm a}$	$0.0330 \pm 0.0040^a$	$0.0408 \pm 0.0010^{\rm d}$	$0.0263 \pm 0.0034^{\rm a}$	-		
T2	$0.0217 \pm 0.0033^{\rm a}$	$0.0280 \pm 0.0017^{\rm c}$	$0.0402 \pm 0.0047^{\rm b}$	$0.0632 \pm 0.0157^{\rm d}$	$0.0254 \pm 0.0016^{c}$		
T3	$0.0217 \pm 0.0033^{\rm a}$	$0.0384 \pm 0.0007^{\rm d}$	$0.0566 \pm 0.0044^{e}$	$0.0190 \pm 0.0077^{c}$	$0.0158 \pm 0.0016^{a}$		
T4	$0.0217 \pm 0.0033^{\rm a}$	$0.0340 \pm 0.0012^{\rm b}$	$0.0465 \pm 0.0037^{c}$	$0.0563 \pm 0.0102^{\rm b}$	$0.0257 \pm 0.0007^{b}$		
C	$0.0217 \pm 0.0033^{\rm a}$	$0.0440 \pm 0.0024^{\circ}$	$0.0504 \pm 0.0004^{\rm a}$	-	-		

Values are mean  $\pm$  standard deviation, n = 3. Values within treatments with different letters (a–d) in a column differ significantly ( $P \le 0.05$ ) with values from higher to lower. T1: Gumghatti 0.6%; T2: Gum-ghatti (0.6% coconut oil (0.3%); T3: Gum-ghatti (0.9%) + jojoba oil (0.3%) and T4: Gum-ghatti (0.6%) + chamomile oil (0.3%) and C: control.

loss than the control group, proving the effectiveness of coating [Table 4]. These results exhibited that the coatings on fruits developed a protective barrier on plum fruit skin that retarded the gaseous exchange and ripening process [36,37]. On the eighth day from the start of the shelf life study, fruits treated with gumghatti + coconut oil displayed a significant rise in lycopene content followed by a gradual decrease. The above trend in the lycopene values aligns with research findings from earlier studies conducted on the coating of tomatoes with the chitosan/zeolite [36], and the combinations of aloe vera and chitosan [37].

## 3.6. Total Phenols and Ascorbic Acid

According to [Table 5], as the storage duration increased, the total phenol content in the fruits decreased. Nonetheless, fruits treated with gumghatti and chamomile oil (T4) exhibited the least decrease in phenolic content. The decrease in total phenolic content by the end

of the storage period was due to the degradation of cell structure as an indication of senescence phenomena during storage [38]. Regarding the ascorbic acid content, it initially increased and then a gradual decrease was noted, with the treated fruits showing higher content compared with untreated fruits ( $p \le 0.05$ ). Fruit sugar's availability, which is a precursor to ascorbic acid production, may have caused an early enhancement in ascorbic acid. However, oxidative ascorbic acid degradation by oxidase may have contributed to a subsequent decline in ascorbic acid [39]. These findings align with earlier studies conducted on tomatoes [38] and sweet cherry fruit [30].

## 3.7. Total Antioxidant Activity

Antioxidants act as free radical scavengers by donating a pair of electrons and neutralizing free radicals, which are oxidizing in nature and harm-causing agents [40]. In the present study, better antioxidant activity was observed in the treated black plums while a rapid loss in antioxidant

activity was noted in the control groups. Anthocyanins were also found to increase, leading to an increase in the total antioxidant activity as a whole. Throughout the storage period, the fruit's total antioxidant activity remained relatively stable, except a slight increase on the eighth day in T3 and T4, followed by a subsequent decrease until the final day of storage [Table 5]. This observation aligns with findings from previous studies in which the papaya fruits were coated with gumghatti and clove oil [11]. However, in the present study, T4 exhibited better antioxidant activity than any other treatments and control on the 16th day of storage.

## 3.8. Polygalactourinase Activity and Pectate Lyase Activity

An increase in PG activity, reaching a peak, followed by a subsequent decline [Table 6], was observed during the storage period. Nonetheless, treated fruits exhibited a slower rise in enzyme activity compared with untreated fruits. Moreover, PL activity also resulted in a significant increase in the duration of storage ( $p \le 0.05$ ). A similar trend of delayed increase in PG activity was reported by Gol *et al.* [41] in strawberry fruit where chitosan was used as edible coating. These coatings provide less accessibility for the above enzymes to act on pectin, leading to reduced pectin solubilization and softening. According to Zhou *et al.* [42], hydrolysis of (1-4) galacturonan linkages of demethylated pectins is catalyzed by PG, followed by the release of shorter chains causing the dissolution of pectins. They also noticed a lower activity of PG in the pears coated with shellac compared with control, thus retaining the firmness of the fruit better.

## 3.9. Amylase Activity

During storage, the amylase activity, responsible for starch break down, showed a significant decrease from the initial to the final day in all groups, while it is slower in treated group compared with control ( $p \le 0.05$ ). Among the treated groups, fruits coated with gumghatti and jojoba oil (T3) exhibited the lowest activity of amylase on the 16th day of storage [Table 6]. In a study performed on edible coating of berry cactus fruit, it was found that sodium caseinate coatings inhibited the amylase activity better than control. This inhibitory activity aids in slower break down of starch, thus resulting in lower glucose levels in blood after consumption [43]. Rahman et al. also found a similar trend in amylase activity pertaining to mangoes stored at three varied temperatures (-5, 4, and  $25^{\circ}$ C); however, it increased with an increase in temperature [44].

## 3.10. Browning Index and PPO Activity

Browning is mainly caused by oxidation of phenolic compounds by an enzyme; the PPO present in fruits. PPO is a key enzyme that activates during senescence, ripening, or stress conditions like membrane damage, resulting in browning of fruit tissues [45]. In the present study, browning phenomena were increased in treated as well as untreated fruits, but this process was found to be comparatively slower in coated black plum fruits than in the control set of fruits [Table 7]. As mentioned above, less intensity of browning was observed in treated fruits due to the coatings that acted as barriers and led to a reduction in the oxygen supply for enzymatic oxidation of contained phenolics [46]. On the fourth day of storage, the control group exhibited higher polyphenol oxidase activity, which is associated with the browning process, compared with the coated fruits [Table 7]. Among the coated fruits, those treated with gumghatti displayed the lowest enzyme activity, while the combination of gumghatti and chamomile oil also contributed to a significant reduction in the enzyme activity comparatively ( $p \le 0.05$ ). The utilization of gumghatti and chamomile oil (T4) resulted in the least browning, comparable to the findings observed in sweet cherry fruits [47].

### 3.11. Shelf Life of Fruits

The control fruits had a shelf life of 7–8 days, whereas the fruits treated with combinations of gumghatti and various oil (T2, T3, and T4) had shelf lives of 15, 15, and 16 days, respectively. It is noteworthy that the combination of gumghatti and chamomile oil significantly doubled the fruits' shelf life ( $p \le 0.05$ ). Furthermore, all treatments not only enhanced the fruits' visual appearance and commercial acceptability but also extended their shelf life by more than 8 days. Thus, the current study reports are in accordance with the results of Vyas *et al.* [48], who reported that polysaccharide-based coatings have a significant role in improving the quality of papaya, thereby extending its shelf life.

## 4. CONCLUSION

Edible coatings are found to be versatile alternatives for enhancing the quality of fresh produce. However, knowledge of the physicochemical and functional characteristics of biopolymers is critical in selecting the appropriate biomaterial specific to the fresh produce. The inherent carrying capacity of the coating materials opens a wide range of opportunities through structural modifications and incorporation of additives like bioactive compounds that might result in better efficiency of the coatings. In the present study, attempt has been made by the application of gumghatti coating, alone and in combination with oils, to enhance the shelf life and postharvest quality characteristics of black plum fruits. The coatings effectively minimized weight and moisture loss, while also enhancing firmness and delaying decay and browning. Furthermore, the treated fruits exhibited elevated levels of essential antioxidants, including lycopene, anthocyanin, and ascorbic acid, leading to delayed ripening compared with the control group. Moreover, the coatings had a notable impact on enzyme activities linked to fruit softening and browning. Among the various coatings, the combination of gumghatti and chamomile oil (T4) demonstrated the highest effectiveness in extending the shelf life of black plum fruits. This combination of chamomile oil and gumghatti can be taken as a promising composite edible coating material for future aspects. This study presents a practical, cost-effective, and user-friendly edible coating that effectively prolongs fruit shelf life even at room temperature.

### 5. ACKNOWLEDGMENTS

The authors wish to express their profound gratitude to the Head, Department of Biosciences, Sardar Patel University, Vallabh Vidyanagar, Anand, Gujarat, for providing the necessary facilities to carry out the experiments.

## 6. AUTHOR CONTRIBUTIONS

Study conception and design: TVRR and PKD; data collection: PKD, JBP and NP; analysis and interpretation of results: TVRR, PRP, and AP; draft manuscript preparation: PKD, TVRR, PRP, AP and NP. The results and final version of the manuscript is approved by all the authors.

## 7. FUNDING

There is no funding to report.

## 8. CONFLICTS OF INTEREST

The authors declare that there is no conflict of interest in this work.

## 9. ETHICAL APPROVAL

This study does not involve experiments on animals or human subjects.

#### 10. DATA AVAILABILITY

All dataset were generated and analyzed in the present study.

# 11. USE OF ARTIFICIAL INTELLIGENCE (AI)-ASSISTED TECHNOLOGY

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

### 12. PUBLISHER'S NOTE

This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

#### REFERENCES

- Hussain SZ, Naseer B, Qadri T, Fatima T, Bhat TA. Plum (Prunus domestica): morphology, taxonomy, composition and health benefits.
  In: Fruits Grown in Highland Regions of the Himalayas. Nutritional and Health Benefits. Cham: Springer International Publishing 2021.
  p. 169-79. https://doi.org/10.1007/978-3-030-75502-7
- Kumar P, Sethi S, Sharma RR, Srivastav M, Singh D, Varghese E. Edible coatings influence the cold-storage life and quality of 'Santa Rosa' plum (*Prunus salicina Lindell*). J Food Sci Technol. 2018;55(6):2344-50. https://doi.org/10.1007/s13197-018-3130-1
- Mitchell FG. Commercial cooling of fruits and vegetables. Berkeley (1972) https://doi.org/10.5962/bhl.title.61202
- Vangdal E, Flatland S, Nordb R. Fruit quality changes during marketing of new plum cultivars (*Prunus domestica* L.). Hort Sci. 2007;34(3):91-5. https://doi.org/10.17221/1892-hortsci
- Sharma S, Sharma R, Pal RK, Singh S. Influence of 1-MCP on compression injury, fruit firmness and quality of Japanese plum cv. Santa Rosa during transportation. Indian J Hort. 2013;70:101-6.
- Mare L, Huysamer M, Dodd MC, Truter AB, Kemp AT, Holcroft DM. Extension of the storage life of plums (*Prunus salicina*) using controlled atmosphere shipping. Acta Hort. 2005;682:1689-96. https://doi.org/10.17660/actahortic.2005.682.225
- Valero D, Diaz-Mula HM, Zapata PJ, Guillen F, Martínez-Romero D, Castillo S, Serrano M. Effects of alginate edible coating on preserving fruit quality in four plum cultivars during postharvest storage. Postharvest Biol Technol. 2013;77:1-6. https://doi.org/10.1016/j. postharvbio.2012.10.011
- Eum HL, Hwang DK, Linke M, Lee SK, Zude M. Influence of edible coating on quality of plum (*Prunus salicina* Lindl. cv. 'Sapphire'). Eur Food Res Technol. 2009;229(3):427-34. https://doi.org/10.1007/ s00217-009-1054-8
- Mondal K, Goud VV, Katiyar V. Effect of waste green algal biomass extract incorporated chitosan-based edible coating on the shelf life and quality attributes of tomato. ACS Food Sci Technol. 2022;2(7):1151-65. https://doi.org/10.1021/acsfoodscitech.2c00174
- Sharma H, Shami V, Samsher, Chaudhary V, Sunil E, Kumar M. Importance of edible coating on fruits and vegetables: a review. J Pharmaco Phytochem. 2019;8:4104-10.
- Joshi AV, Baraiya NS, Vyas PB, Rao TVR. Gumghatti based edible coating emulsion with an additive of clove oil improves the storage life and maintains the quality of papaya (Carica papaya L., cv. Madhu

- bindu). Int J Curr Microbiol Appl Sci. 2017;6(5):160-74. https://doi.org/10.20546/ijcmas.2017.605.019
- Mondal MF. Production and storage of fruits (in Bangla). Mymensingh: BAU Campus; 2000.
- Horwitz W. Official methods of analysis. Vol. 222. Washington (DC): Association of Official Analytical Chemists; 1975.
- 14. Sadasivam S, Manickam A. Biochemical methods for agricultural sciences. Wiley Eastern Ltd., New Delhi; 1992.
- Eissa HAA. Effect of Chitosan coating on shelf life and quality of fresh-cut mushroom. J Food Qual. 2007;30(5):623-45. https://doi. org/10.1111/j.1745-4557.2007.00147.x
- Lees DH, Francis FJ. Standardization of pigment analyses in cranberries1. HortScience. 1972;7(1):83-4. https://doi.org/10.21273/ hortsci.7.1.83
- Larrauri JA, Sanchez-Moreno C, Saura-Calixto F. Effect of temperature on the free radical scavenging capacity of extracts from red and white grape pomace peels. J Agri Food Chem. 1998;46(7):2694-7. https://doi.org/10.1021/jf980017p
- McDonald S, Prenzler PD, Antolovich M, Robards K. Phenolic content and antioxidant activity of olive extracts. Food Chem. 2001;73(1):73-84. https://doi.org/10.1016/s0308-8146(00)00288-0
- Roe JH, Kuether CA. The determination of ascorbic acid in whole blood and urine through the 2,4-dinitrophenylhydrazine derivative of dehydroascorbic acid. J Biol Chem. 1943;147(2):399-407. https:// doi.org/10.1016/s0021-9258(18)72395-8
- Wang LZ, Liu L, Holmes J, Kerry JF, Kerry JP. Assessment of filmforming potential and properties of protein and polysaccharide-based biopolymer films. Inter J Food Sci Technol. 2007;42(9):1128-38. https://doi.org/10.1111/j.1365-2621.2006.01440.x
- Ouattara HG, Reverchon S, Niamke SL, Nasser W. Biochemical properties of pectate lyases produced by three different bacillus strains isolated from fermenting cocoa beans and characterization of their cloned genes. Appl Env Microbiol. 2010;76(15):5214-20. https://doi.org/10.1128/aem.00705-10
- Pathak N, Sanwal GG. Multiple forms of polygalacturonase from banana fruits. Phytochem. 1998;48(2):249-55. https://doi. org/10.1016/s0031-9422(98)00005-3
- Xing Y, Li X, Xu Q, Yun J, Lu Y, Tang Y. Effects of chitosan coating enriched with cinnamon oil on qualitative properties of sweet pepper (Capsicum annuum L.). Food Chem. 2011;124(4):1443-50. https:// doi.org/10.1016/j.foodchem.2010.07.105
- Bliss CI. Statistical methods for research in the natural sciences. In: Statistics in Biology. London: McGraw Hill Book Company; 1967. p. 558.
- Rojas-Argudo C, Del Río MA, Perez-Gago MB. Development and optimization of locust bean gum (LBG) based edible coatings for postharvest storage of Fortune mandarins. Postharvest Biol Technol. 2009;52:227-34.
- Kittur FS, Saroja N, Habibunnisa, Tharanathan RN. Polysaccharide based composite coating formulations for shelf-life extension of fresh banana and mango. Eur Food Res Technol. 2001;213:306-11.
- 27. Thomas SA, Molina, E-Bosques, Stolik S, Sanchez F. Composite oil coating preserves the quality of pineapple fruits. J Physiq. 2005;125:889-92.
- Zapata PJ, Guille F, Martinez-Romero D, Castillo S, Valero D, Serrano M. Use of alginate or zein as edible coatings to delay postharvest ripening process and to maintain tomato (Solanum lycopersicon Mill) quality. J Sci Food Agric. 2008;88:1287-93.
- 29. Salunkhe DK, Boun HR, Reddy NR. Storage processing and nutritional quality of fruits and vegetables, Vol. 1. Fresh Fruits and Vegetables. Bostan: CRC Press Inc; 1991.
- 30. Mahfoudhi N, Hamdi S. Use of almond gum and gum arabic as novel edible coating to delay postharvest ripening and to maintain

- sweet cherry (*Prunus avium*) quality during storage. J Food Process Preserv. 2015;39(6):1499-508.
- 31. Pandey SK, Joshua JE, Bisen, Abhay. Influence of gamma irradiation, growth retardants and coatings on the shelf life of winter guava fruits (*Psidium guajava* L.). J Food Sci. Technol. 2012;49(6):753-9.
- 32. Zhang W, Guo M, Yang W, Liu Y, Wang Y, Chen G. The role of cell wall polysaccharides disassembly and enzyme activity changes in the softening process of hami melon (*Cucumis melo* L.) Foods. 2022;11(6):841.
- Debeaufort FJ, Quezada-Gallo A, Voilley A. Edible films and coatings: tomorrow's packaging: a review. Crit Rev Food Sci Nutr. 1998;38:299-313.
- Zam W. Effect of alginate and chitosan edible coating enriched with olive leaves extract on the shelf life of sweet cherries (*Prunus avium L.*) J Food Qual. 2019;2019:8192964. https://doi. org/10.1155/2019/8192964
- 35. Usenik V, Kastelec D, Veberič R, Štampar F. Quality changes during ripening of plums (*Prunus domestica* L.) Food Chem. 2008;111(4):830-6.
- Garca M, Casariego A, Roblejo R. Effect of edible chitosan/ zeolite coating on tomatoes quality during refrigerated storage. Emirates J Food Agri. 2014;26(3):238. https://doi.org/10.9755/ejfa. v26i3.16620
- Khatri D, Panigrahi J, Prajapati A, Bariya H. Attributes of Aloe vera gel and chitosan treatments on the quality and biochemical traits of post-harvest tomatoes. Sci Horticult. 2020;259:108837.
- Dávila-Aviña JE, Villa-Rodríguez JA, Villegas-Ochoa MA, Tortoledo-Ortiz O, Olivas GI, Ayala-Zavala JF, et al. Effect of edible coatings on bioactive compounds and antioxidant capacity of tomatoes at different maturity stages. J Food Sci Technol. 2012;51(10):2706-12. https://doi.org/10.1007/s13197-012-0771-3
- Chawla S, Devi R, Jain V. Changes in physicochemical characteristics of guava fruits due to chitosan and calcium chloride treatments during storage. J Pharmacognosy Phytochem. 2018;7(3):1035-44.
- Jhalegar MDJ, Sharma RR, Pal RK, Rana V. Effect of postharvest treatments with polyamines on physiological and biochemical

- attributes of kiwifruit (*Actinidia deliciosa*) cv. Allison. Fruits. 2012;67(1):13-22.
- Gol NB, Patel RP, Rao TVR. Improvement of quality and shelf life of strawberry with edible coatings enriched with chitosan. Postharvest Biol Technol. 2013;85:185-95.
- 42. Zhou, R, Li Y, Yan L, Xie J. Effect of edible coatings on enzymes, cell-membrane integrity and cell-wall constituents in relation to brittleness and firmness of Huanghua pears (*Pyrus pyrifolia* Nakai, cv. Huanghua) during storage. Food Chem. 2011;124:569-75.
- 43. Correa-Betanzo J, Jacob JK, Perez-Perez C, Paliyath G. Effect of a sodium caseinate edible coating on berry cactus fruit (Myrtillocactus geometrizans) phytochemicals. Food Res Int. 2011;44(7):1897-904.
- 44. Rahman MM, Rahman MM, Absar N, Ahsan MA. Correlation of carbohydrate content with the changes in amylase, invertase and -galactosidase activity of ripe mango pulp during storage under different temperatures. Bangladesh J Sci Indust Res. 2011;46(4):443-6.
- 45. Mayer AM. Polyphenol oxidase and peroxidase in plants recent progress. Phytochemistry. 1987;26:11-20.
- Panahirad S, Naghshiband-Hassani R, Mahna N. Pectin-based edible coating preserves antioxidative capacity of plum fruit during shelf life. Food Sci Technol Int. 2020;26(7):583-92.
- Díaz-Mula, H. M., Serrano, M., Valero, D. Alginate coatings preserve fruit quality and bioactive compounds during storage of sweet cherry fruit. Food Bioprocess Technol. 2011;5(8):2990-7. https://doi. org/10.1007/s11947-011-0599-2
- Vyas PB, Gol NB, Rao TVR. Postharvest quality maintenance of papaya fruit using polysaccharide-based edible coatings. Int J Fruit Sci. 2013;14(1):81-94. https://doi.org/10.1080/15538362.2013.80

#### How to cite this article:

Rao TVR, Dave PK, Payal A, Pandya JB, Patel PR, Pareek N. Gumghatti-based composite coating improves postharvest quality and nutritional value of black plums. J App Biol Biotech. 2025;13(1):75-82. DOI: 10.7324/JABB.2024.159375